



Atomic-Scale Evidence for the Nucleation Sites of Multilayer Graphene and Its Dynamic Evolution to a Carbon Nanotube

Zhang, Lili; He, Maoshuai; Wagner, Jakob Birkedal

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Zhang, L., He, M., & Wagner, J. B. (2015). *Atomic-Scale Evidence for the Nucleation Sites of Multilayer Graphene and Its Dynamic Evolution to a Carbon Nanotube*. Abstract from Microscopy Conference 2015, Göttingen, Germany.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Atomic-Scale Evidence for the Nucleation Sites of Multilayer Graphene and Its Dynamic Evolution to a Carbon Nanotube

Lili Zhang,¹ Maoshuai He,² Jakob B. Wagner¹

¹Technical University of Denmark, Center for Electron Nanoscopy, Fysikvej 307, 2800 Kgs. Lyngby, Denmark

²Laboratoire d'Étude des Microstructures, ONERA-CNRS, BP 72, 92322 Châtillon CEDEX, France
Phone: +45-5269-6573
E-mail: lilizha@dtu.dk

In situ environmental transmission electron microscopy (ETEM) is of great importance to provide direct experimental evidence of structural dynamics of nanomaterials in a reactive gas at elevated temperature.^{1,2} Here, the catalytic nucleation of carbon nanostructures from nanoparticles are recorded with aberration-corrected high-resolution TEM studies. Figure 1. shows the surface-bounded graphene on a certain facet of the particle stacks layer by layer, meanwhile, the disordered region inside the particle expands deeply along the interface of particle and support, suggesting a bulk diffusion path of carbon. However, the appearance of the gas-absorbed interface on the other facet of the particle allows assembling a graphene layer on its partial surface via surface diffusion. The evolution of the particle morphology then results in an increased number of layers and a tube-like structure. Overall, this work provides an improved understanding of nucleation mechanism of graphene and carbon nanotubes.

(1) Hansen, T. W.; Wagner, J. B. *ACS Catal.* **2014**, 4, 1673.

(2) He, M.; Kauppinen, E. I. et al. *Sci. Rep.* **2013**, 3.




Figure 1. A series of high-resolution TEM images showing the nucleation processes of multilayer graphene. (a-c) A stack of graphene on one facet of the particle. (d) A graphene layer assembles on the other facet of the particle. Fast fourier transforms (FFTs) from the squares showing the expanded disordered region inside the particle.